arXiv:1112.0112v1 [math-ph] 1 Dec 2011

Zero Energy Bound States in Many–Particle Systems

Dmitry K. Gridnev†

FIAS, Ruth-Moufang-Straße 1, D–60438 Frankfurt am Main, Germany

E-mail: `gridnev@fias.uni-frankfurt.de`

Abstract. It is proved that the eigenvalues in the N–particle system are absorbed at zero energy threshold, if none of the subsystems has a bound state with $E \leq 0$ and none of the particle pairs has a zero energy resonance. The pair potentials are allowed to take both signs.

1. Introduction

In [1] it was proved that the 3–body system, which is at the 3–body coupling constant threshold, has a square integrable state at zero energy if none of the 2–body subsystems is bound or has a zero energy resonance. The condition on the absence of 2–body zero energy resonances is essential, that is the 3–body ground state at zero energy can be at most a resonance and not an L^2 state if at least one pair of particles has a zero energy resonance [1]. One of the restrictions on pair potentials in [1] was their being non–positive. The aim of the present paper is to generalize the result of [1] to the case of many particles and get rid of the restriction on the sign of pair potentials. The main result is expressed in Theorems 1, 2, which state that the eigenvalues in the N–particle system are absorbed at zero energy threshold, if none of the subsystems has a bound state with $E \leq 0$ and none of the particle pairs has a zero energy resonance. Throughout the paper we use the following operator notation. $A \geq 0$ means that $(f, Af) \geq 0$ for all $f \in D(A)$ and $A \ngeq 0$ means that there exists $f_0 \in D(A)$ such that $(f_0, Af_0) < 0$.

We consider the N–particle Schrödinger operator

$$H(\lambda) = H_0 + \lambda \sum_{1\leq i<j\leq N} V_{ij}(r_i - r_j), \tag{1}$$

where $\lambda > 0$ is a coupling constant, H_0 is a kinetic energy operator with the centre of mass removed, $r_i \in \mathbb{R}^3$ are particle position vectors, the pair potentials are real and $V_{ij} \in L^2(\mathbb{R}^3) \cap L^1(\mathbb{R}^3)$. The operator $H(\lambda)$ is self–adjoint on $D(H_0) \subset L^2(\mathbb{R}^{3N-3})$, the set of relative coordinates in $\mathbb{R}^{3N-3}$ we shall denote as ξ. Throughout the paper we shall assume that

$$\sigma_{ess}(H(\lambda)) = [0, \infty), \tag{2}$$

---

† On leave from: Institute of Physics, St. Petersburg State University, Ulyanovskaya 1, 198504 Russia

which, of course, restricts possible values of λ. Here we shall extensively use the term *critical coupling*. In the literature one finds several related definitions: critically bound [2], critical coupling [3], coupling constant threshold in [4, 5], virtual level at the threshold [6], etc. To avoid possible confusion we list three of the most popular definitions and indicate the relations between them.

Definition 1. *$H(\lambda)$ is at critical coupling if $H(\lambda) \geq 0$ and $H(\lambda) + \epsilon \sum_{i<j} V_{ij} \ngeq 0$ for any $\epsilon > 0$.*

In the terminology of [4, 5] Def. 1 implies that $H(\lambda)$ is at the coupling constant threshold. The next definition due to scaling arguments is fully equivalent to Def. 1

Definition 2. *$H(\lambda)$ is at critical coupling if $H(\lambda) \geq 0$ and $H(\lambda) - (1-\epsilon)H_0 \ngeq 0$ for any $0 < \epsilon < 1$.*

So under the term *critical coupling* we shall mean any of those. The next definition can be found, for example, in [6]

Definition 3. *$H(\lambda)$ is said to have a virtual level at zero energy if $H(\lambda) \geq 0$ and $H(\lambda) - \epsilon V_R \ngeq 0$ for any $\epsilon > 0$, where $V_R := 1/(1+|\xi|^2)$.*

In the case of $N = 2$ it is easy to show that all three definitions are equivalent to the definition of a two–particle zero energy resonance, c.f. [7, 8]. Note, that, in general, for $N \geq 3$ the Defs. 1–2 and Def. 3 are not equivalent. The difference lies in the fact that the perturbation in Def. 3 does not move the lower bound of the essential spectrum, since V_R is a relatively H_0–compact perturbation, contrary to the perturbations in Defs. 1–2, where the lower bound of the essential spectrum can be moved, if some of the subsystems are at critical coupling.

Proposition 1. *A system of N particles is at critical coupling if it has a virtual level at zero energy.*

Proof. Assume by contradiction that the system is not at critical coupling. Then there must exist $\epsilon_0 > 0$ such that $H - \epsilon_0 H_0 \geq 0$. By the Courant identity [9, 10] there exists $\kappa > 0$ such that $H_0 - \kappa V_R \geq 0$. Hence,

$$H - \epsilon_0 \kappa V_R = H - \epsilon_0 H_0 + \epsilon_0 (H_0 - \kappa V_R) \geq 0, \tag{3}$$

which means that the system does not have a virtual level at zero energy. □

As already mentioned the converse of Proposition 1 is in general not true. Note, however, that that if a system has a zero energy bound state then it automatically has a virtual level at zero energy.

2. Main Result

For the formulation of Theorem 1 we need to impose the following requirement.

- R1 There exists a sequence of coupling constants $\lambda_n \in \mathbb{R}_+$ such that $\lim_{n\to\infty} \lambda_n = \lambda_{cr} \in \mathbb{R}_+$, and $H(\lambda_n)\psi_n = E_n\psi_n$, where $\psi_n \in D(H_0)$, $\|\psi_n\| = 1$, $E_n < 0$, $\lim_{n\to\infty} E_n = 0$.

Further in this section we shall prove the following

Theorem 1. *Suppose* $H(\lambda)$ *defined in (1) for* $N \geq 3$ *satisfies R1,* $H(\lambda_n)$ *and* $H(\lambda_{cr})$ *have no subsystems, which have a bound state with* $E \leq 0$*, and no particle pairs at critical coupling. Then there exists normalized* $\psi_0 \in D(H_0)$ *such that* $H(\lambda_{cr})\psi_0 = 0$*.*

The next statement can be considered as a corollary to Theorem 1.

Theorem 2. *Suppose that* $N \geq 3$ *and* $H(\lambda_{cr})$ *is at critical coupling. Suppose also that* $H(\lambda_{cr})$ *has no subsystems, which have a bound state with* $E \leq 0$*, and no particle pairs at critical coupling. Then there exists normalized* $\psi_0 \in D(H_0)$ *such that* $H(\lambda_{cr})\psi_0 = 0$*.*

Proof. Let us assume that none of the subsystems is at critical coupling. On one hand, from the HVZ theorem [11, 10] it follows that there exists $\epsilon_0 > 0$ such that for $\lambda_n = \lambda_{cr}(1 + \epsilon_0/n)$ and $n = 1, 2, \dots$ we have $\inf \sigma_{ess} H(\lambda_n) = 0$. We also choose ϵ_0 small enough to guarantee that $H(\lambda_n)$ has no subsystems that are either bound or at critical coupling. On the other hand, $H(\lambda_n) \ngeq 0$. Therefore, there are $\psi_n \in D(H_0)$ such that $H(\lambda_n)\psi_n = E_n\psi_n$, where $E_n < 0$, $\|\psi_n\| = 1$ and $E_n \to 0$. Now the statement follows from Theorem 1. It remains to get rid of the assumption that there is no subsystems at critical coupling. If there would be such then it is always possible to pass to the corresponding subsystem (call it $\mathcal{S}$), which has no subsystems at critical coupling. In such case by the above analysis $\mathcal{S}$ must have a bound state with $E = 0$, which is in contradiction with the theorem conditions. □

Following [1] let us introduce the operator $B_{\tau_1\tau_2}(z)$, where $1 \leq \tau_1 < \tau_2 \leq N$. We construct $B_{12}(z)$, for other particle pairs the construction is analogous.

We use Jacobi coordinates [12] $\xi = (x, y_1, y_2, \dots, y_{N-2})$, where $x, y_i \in \mathbb{R}^3$. We set $x = \alpha^{-1}(r_2 - r_1)$ and $y_1 = (\sqrt{2M_{12}}/\hbar)[r_3 - m_1/(m_1 + m_2)r_1 - m_2/(m_1 + m_2)r_2]$, where $\alpha := \hbar/\sqrt{2\mu_{12}}$, $M_{12} := (m_1 + m_2)m_3/(m_1 + m_2 + m_3)$ and $\mu_{ik} := m_i m_k/(m_i + m_k)$ is the reduced mass. For $N = 4$ this choice of coordinates is illustrated in Fig. 1 (Left). The coordinate $y_i \in \mathbb{R}^3$ is proportional to the vector pointing from the centre of mass of the particles $[1, 2, \dots, i + 1]$ to the particle $i + 2$, and the scale is set to make the kinetic energy operator take the form

$$H_0 = -\Delta_x - \sum_i \Delta_{y_i}. \tag{4}$$

Let $\mathcal{F}_{12}$ denote the partial Fourier transform in $L^2(\mathbb{R}^{3N-3})$ acting as follows

$$\hat{f}(x, p_y) = \mathcal{F}_{12} f = \frac{1}{(2\pi)^{(3N-6)/2}} \int d^{3N-6}y \ \ e^{-ip_y\cdot\, y} f(x, y), \tag{5}$$

where $y = (y_1, \dots, y_{N-2})$, $p_y = (p_{y_1}, p_{y_2}, \dots, p_{y_{N-2}}) \in \mathbb{R}^{3N-6}$. Then $B_{12}(z)$ is defined through

$$B_{12}(z) = 1 + z + \mathcal{F}_{12}^{-1} t(p_y) \mathcal{F}_{12}, \tag{6}$$

where

$$t(p_y) = \left(\sqrt{|p_y|} - 1\right) \chi_{\{p_y |\ |p_y| \leq 1\}}, \tag{7}$$

$|p_y| = (\sum_i p_{y_i}^2)^{1/2}$ and χ_Ω denotes the characteristic function of the set Ω. Let us transform the coordinates through $\tilde{y}_i = \sum_k T_{ik} y_k$, where T_{ik} is any orthogonal $(N-2) \times (N-2)$ matrix. It is easy to check that the construction of $B_{12}(z)$ is invariant with respect to these coordinate transformations. That is

$$B_{12}(z) = 1 + z + \tilde{\mathcal{F}}_{12}^{-1} t(\tilde{p}_y) \tilde{\mathcal{F}}_{12}, \tag{8}$$

where $\tilde{\mathcal{F}}_{12}$ is defined through

$$\hat{f}(x, \tilde{p}_y) = \tilde{\mathcal{F}}_{12} f = \frac{1}{(2\pi)^{(3N-6)/2}} \int d^{3N-6}\tilde{y}\ \ e^{-i\tilde{p}_y \cdot\, \tilde{y}} f(x, \tilde{y}). \tag{9}$$

Similarly, one defines $B_{\tau_1\tau_2}(z)$ for all particle pairs. $B_{\tau_1\tau_2}(z)$ and $B^{-1}_{\tau_1\tau_2}(z)$ are analytic on $\mathrm{Re}\, z > 0$.

Proof of Theorem 1. By contradiction, let us assume that the zero energy bound state does not exist. Then by Theorem 1 in [1] ψ_n totally spreads and $\psi_n \overset{w}{\to} 0$ (for the definition of spreading see [1]). Let $\tau = 1, 2, \ldots, N(N-1)/2$ for $N \geq 4$ label all particle pairs and $\tau_1 < \tau_2$ label the particle numbers entering the pair τ. We shall denote $v_\tau := V_{\tau_1\tau_2}$. It is helpful to split v_τ into positive and negative parts $v_\tau = (v_\tau)_+ - (v_\tau)_-$, where $(v_\tau)_+ := \max[0, v_\tau]$ and $(v_\tau)_- := \max[0, -v_\tau]$. On one hand, the Schrödinger equation for ψ_n reads

$$\left(H_0 + \lambda_n U_+ + k_n^2\right)\psi_n = \lambda_n \sum_\tau \sqrt{(v_\tau)_-}(\sqrt{(v_\tau)_-}\psi_n), \tag{10}$$

where we set

$$U_+ := \sum_\tau (v_\tau)_+. \tag{11}$$

Acting on the last equation with an inverse operator gives

$$\psi_n = \lambda_n \sum_\tau \left(H_0 + \lambda_n U_+ + k_n^2\right)^{-1} \sqrt{(v_\tau)_-}\left(\sqrt{(v_\tau)_-}\psi_n\right). \tag{12}$$

On the other hand, we can rearrange the terms in the Schrödinger equation as follows

$$\left[H_0 + \lambda_n (v_\tau)_+ + k_n^2\right]\psi_n = \lambda_n (v_\tau)_- \psi_n - \lambda_n \sum_{\delta \neq \tau} v_\delta \psi_n, \tag{13}$$

where index δ runs through all particle pairs. This gives us

$$\psi_n = \lambda_n \left[H_0 + \lambda_n (v_\tau)_+ + k_n^2\right]^{-1} (v_\tau)_- \psi_n - \lambda_n \sum_{\delta \neq \tau} \left[H_0 + \lambda_n (v_\tau)_+ + k_n^2\right]^{-1} v_\delta \psi_n. \tag{14}$$

Using (14) we get the following expression for the last term in brackets in (12)

$$\sqrt{(v_\tau)_-}\psi(\lambda_n) = -\lambda_n \sum_{\delta\neq\tau} \Big\{ 1 - \lambda_n \sqrt{(v_\tau)_-} \Big[H_0 + \lambda_n (v_\tau)_+ + k_n^2 \Big]^{-1} \sqrt{(v_\tau)_-} \Big\}^{-1} \sqrt{(v_\tau)_-} \Big[H_0 + \lambda_n (v_\tau)_+ + k_n^2 \Big]^{-1} v_\delta \psi_n. \tag{15}$$

That the inverse of the operator in curly brackets makes sense would be shown in Lemma 1 below. Substituting (15) into (12) yields the equation

$$\psi_n = -\lambda_n^2 \sum_\tau \sum_{\delta\neq\tau} \Big(H_0 + \lambda_n U_+ + k_n^2 \Big)^{-1} \sqrt{(v_\tau)_-} \Big\{ 1 - \lambda_n \sqrt{(v_\tau)_-} \Big[H_0 + \lambda_n (v_\tau)_+ + k_n^2 \Big]^{-1} \sqrt{(v_\tau)_-} \Big\}^{-1} \sqrt{(v_\tau)_-} \Big[H_0 + \lambda_n (v_\tau)_+ + k_n^2 \Big]^{-1} v_\delta \psi_n \tag{16}$$

All operators under the sum except v_δ are positivity preserving, see [13, 11, 10]. The inverse of the operator in curly brackets being positivity preserving can be seen from its expansion in von Neumann series, see and Lemma 12 in [1] and Lemma 1 of this paper. Thus we can transform (16) into the following inequality

$$|\psi_n| \leq \lambda_n^2 \sum_\tau \sum_{\delta\neq\tau} \Big(H_0 + \lambda_n U_+ + k_n^2 \Big)^{-1} \sqrt{(v_\tau)_-} \Big\{ 1 - \lambda_n \sqrt{(v_\tau)_-} \Big[H_0 + \lambda_n (v_\tau)_+ + k_n^2 \Big]^{-1} \sqrt{(v_\tau)_-} \Big\}^{-1} \sqrt{(v_\tau)_-} \Big[H_0 + \lambda_n (v_\tau)_+ + k_n^2 \Big]^{-1} |v_\delta| |\psi_n|. \tag{17}$$

Note that $\Big(H_0 + \lambda_n U_+ + k_n^2\Big)^{-1}$ and $\Big[H_0 + \lambda_n (v_\tau)_+ + k_n^2\Big]^{-1}$ are integral operators, see [13], and positivity preserving operators, see, for example, [11] (Example 3 from Sec. IX.7 in vol. 2 and Theorem XIII.44 in vol. 4). By resolvent identities

$$\Big(H_0 + k_n^2\Big)^{-1} - \Big(H_0 + \lambda_n U_+ + k_n^2\Big)^{-1} = \lambda_n \Big(H_0 + \lambda_n U_+ + k_n^2\Big)^{-1} U_+ \Big(H_0 + k_n^2\Big)^{-1}. \tag{18}$$

$$\Big(H_0 + k_n^2\Big)^{-1} - \Big(H_0 + \lambda_n (v_\tau)_+ + k_n^2\Big)^{-1} = \lambda_n \Big(H_0 + \lambda_n (v_\tau)_+ + k_n^2\Big)^{-1} (v_\tau)_+ \Big(H_0 + k_n^2\Big)^{-1} \tag{19}$$

the differences on the lhs of (18)–(19) are positivity preserving operators. Therefore, we can rewrite (17) as

$$|\psi_n| \leq \lambda_n^2 \sum_\tau \sum_{\delta\neq\tau} \Big(H_0 + k_n^2 \Big)^{-1} \sqrt{(v_\tau)_-} \Big\{ 1 - \lambda_n \sqrt{(v_\tau)_-} \Big[H_0 + \lambda_n (v_\tau)_+ + k_n^2 \Big]^{-1} \sqrt{(v_\tau)_-} \Big\}^{-1} \sqrt{(v_\tau)_-} \Big[H_0 + k_n^2 \Big]^{-1} |v_\delta| |\psi_n|. \tag{20}$$

We use the notation $B_\tau(z) \equiv B_{\tau_1\tau_2}(z)$, where $B_{\tau_1\tau_2}(z)$ was defined above. Inserting into (20) the identity $1 = B_\tau(k_n) B_\tau^{-1}(k_n)$ and using that $[B_\tau(k_n), H_0] = 0$ and $[B_\tau(k_n), (v_\tau)_\pm] = 0$ we obtain

$$|\psi_n| \leq \lambda_n^2 \sum_\tau \sum_{\delta\neq\tau} A_\tau(k_n) R_\tau(k_n) D_{\tau;\delta}(k_n) \sqrt{|v_\delta|} |\psi_n|, \tag{21}$$



Figure 1. Illustration to the choice of Jacobi coordinates for $N = 4$. Left: y_1 points in the direction from the centre of mass of particles [1,2] to the particle 3 and y_2 points in the direction from the centre of mass of particles [1,2,3] (symbolized by a square) to the particle 4. Right: $\tilde{y}_2$ points in the direction from the centre of mass of particles [1,2] to the centre of mass of particles [3,4]. The coordinates' scales are set to make Eq. (4) hold.

where we defined the operators

$$A_\tau(k_n) := \Big(H_0 + k_n^2\Big)^{-1} \sqrt{(v_\tau)_-}B_\tau(k_n), \tag{22}$$

$$R_\tau(k_n) := \Big\{1 - \lambda_n \sqrt{(v_\tau)_-}\Big[H_0 + \lambda_n (v_\tau)_+ + k_n^2\Big]^{-1} \sqrt{(v_\tau)_-}\Big\}^{-1}, \tag{23}$$

$$D_{\tau;\delta}(k_n) := \sqrt{(v_\tau)_-}\Big[H_0 + k_n^2\Big]^{-1} B_\tau^{-1}(k_n)\sqrt{|v_\delta|} \qquad (\tau \neq \delta). \tag{24}$$

Note that $H(\lambda_{cr})$ satisfies the conditions of Theorem 3 in Appendix. It is easy to see that $(\psi_n, H(\lambda_{cr})\psi_n) \to 0$, where ψ_n totally spreads. Hence, by Theorem 3 we have $\|\sqrt{|v_\delta|}\psi_n\| = (\psi_n, |v_\delta|\psi_n) \to 0$. Applying Lemmas 1, 2 to the rhs of (21) tells us that it goes to zero in norm, which is a contradiction, since $\|\psi_n\| = 1$ by R1. □

Lemma 1. *The operators $A_\tau(k_n), R_\tau(k_n)$ given by (22)–(23) are uniformly norm–bounded.*

Proof. Without loosing generality we can consider the pair $\tau = (1, 2)$, where $\tau_1 = 1$ and $\tau_2 = 2$. The proof that $\|A_{12}(k_n)\|$ is uniformly bounded follows the same pattern as the proof of Lemma 6 in [1] and we omit it here. The proof for $R_\tau(k_n)$ uses the Birman–Schwinger principle in the form suggested in [14]. Note that for self–adjoint operators $\mathcal{A}, \mathcal{B} \geq 0$, where $\mathcal{A}^{-1}$ and $\mathcal{A}^{-1/2}\mathcal{B}^{1/2}$ are bounded, one has

$$\|\mathcal{A}^{-1/2}\mathcal{B}\mathcal{A}^{-1/2}\| = \|\mathcal{B}^{1/2}\mathcal{A}^{-1}\mathcal{B}^{1/2}\|, \tag{25}$$

which follows from $\|\mathcal{C}^\dagger\mathcal{C}\| = \|\mathcal{C}\mathcal{C}^\dagger\|$ for any bounded $\mathcal{C}$, see f. e. [15]. Due to conditions of Theorem 1 there exists $1 > \omega' > 0$ (independent of n) such that $(1-\omega')H_0 + \lambda_n(v_\tau) \geq 0$, or, equivalently

$$H_0 + \lambda_n(v_\tau) \geq \omega' H_0. \tag{26}$$

By standard estimates there must exist $\gamma_0 > 0$ such that $H_0 - \gamma_0\lambda_n(v_\tau)_- \geq 0$ for all n. Together with (26) this means that there exists $\omega > 0$ independent of n such that

$$H_0 + \lambda_n(v_\tau) - \lambda_n\omega(v_\tau)_- \geq 0. \tag{27}$$

To use the identity (25) let us set

$$\mathcal{A} := H_0 + \lambda_n (v_\tau)_+ + k_n^2, \tag{28}$$

$$\mathcal{B} := \lambda_n (v_\tau)_-. \tag{29}$$

Because $\mathcal{A} - (1+\omega)\mathcal{B} \geq 0$ for any $\phi \in D(H_0)$ we have

$$(\phi, [\mathcal{A} - (1+\omega)\mathcal{B}]\phi) = (\tilde{\phi}, \{1 - (1+\omega)\mathcal{A}^{-1/2}\mathcal{B}\mathcal{A}^{-1/2}\}\tilde{\phi}) \geq 0, \tag{30}$$

where $\tilde{\phi} := \mathcal{A}^{1/2}\phi$. For $\phi \in D(H_0)$ the functions $\tilde{\phi}$ span a dense set in $L^2(\mathbb{R}^{3N-3})$ since $D(H_0)$ is dense and $\mathcal{A}^{-1/2}$ is bounded. Hence, $(1+\omega)\mathcal{A}^{-1/2}\mathcal{B}\mathcal{A}^{-1/2} \leq 1$ and $\|\mathcal{A}^{-1/2}\mathcal{B}\mathcal{A}^{-1/2}\| \leq 1/(1+\omega)$. By identity (25) we obtain

$$\|\lambda_n \sqrt{(v_\tau)_-}\Big[H_0 + \lambda_n (v_\tau)_+ + k_n^2\Big]^{-1} \sqrt{(v_\tau)_-}\| = \|\mathcal{B}^{1/2}\mathcal{A}^{-1}\mathcal{B}^{1/2}\| \leq 1/(1+\omega), \tag{31}$$

which means that $R_\tau(k_n)$ in (24) is correctly defined and uniformly norm–bounded. □

Lemma 2. *The operators $D_{\tau;\delta}(k_n)$ given by (24) are uniformly norm–bounded.*

Proof. Again it suffices to consider the pair $\tau = (1,2)$, where $\tau_1 = 1$ and $\tau_2 = 2$. We split $D_{12;\delta}(k_n)$ as follows

$$D_{12;\delta}(k_n) = D^{(1)}_{12;\delta}(k_n) + D^{(2)}_{12;\delta}(k_n), \tag{32}$$

$$D^{(1)}_{12;\delta}(k_n) := \sqrt{(v_{12})_-}\Big[H_0 + k_n^2\Big]^{-1}\left\{B_{12}^{-1}(k_n) - \frac{1}{k_n+1}\right\}\sqrt{|v_\delta|}, \tag{33}$$

$$D^{(2)}_{12;\delta}(k_n) := (k_n+1)^{-1}\sqrt{(v_{12})_-}\Big[H_0 + k_n^2\Big]^{-1}\sqrt{|v_\delta|}. \tag{34}$$

For the operator in (34) we get (see Eqs. (43)–(44) in [1])

$$\left\|D^{(2)}_{12;\delta}(k_n)\right\| \leq \left\|\sqrt{(v_{12})_-}\Big[H_0 + k_n^2\Big]^{-1}\sqrt{(v_{12})_-}\right\|^{1/2}\left\|\sqrt{|v_\delta|}\Big[H_0 + k_n^2\Big]^{-1}\sqrt{|v_\delta|}\right\|^{1/2}, \tag{35}$$

where both norms in the product are uniformly bounded (this can be easily shown after making an appropriate Fourier transform). It remains to prove that $D^{(1)}_{12;\delta}(k_n)$ is uniformly norm–bounded. Let us first consider two cases: (a) $\delta_1 = 2$, $\delta_2 = 3$ and (b) $\delta_1 = 3$, $\delta_2 = 4$. The proof for the case (a) almost repeats the one in Lemma 9 in [1]. Indeed, we need to show that $\|\mathcal{K}_n\|$ is uniformly bounded, where

$$\mathcal{K}_n = \mathcal{F}_{12} D^{(1)}_{12;23}(k_n)\mathcal{F}_{12}^{-1}. \tag{36}$$

For convenience we denote $p_{y_r} := (p_{y_2}, p_{y_3}, \ldots, p_{y_{N-2}}) \in \mathbb{R}^{3N-9}$. The integral operator $\mathcal{K}_n$ acts on $\phi(x, p_{y_1}, p_{y_r}) \in L^2(\mathbb{R}^{3N-3})$ as follows

$$\mathcal{K}_n\phi(x, p_{y_1}, p_{y_r}) = \int d^3x'\, d^3p'_{y_1}\; K_n(x, x', p_{y_1}, p'_{y_1}; p_{y_r})\phi(x', p'_{y_1}, p_{y_r}), \tag{37}$$

where the integral kernel has the form [1]

$$K_n(x, x', p_{y_1}, p'_{y_1}; p_{y_r}) = \frac{1}{2^{7/2}\pi^{5/2}\gamma^3}\left[\frac{1}{k_n + 1 + t(p_y)} - \frac{1}{k_n+1}\right]\Big|(V_{12})_-(\alpha x)\Big|^{1/2}$$
$$\times \frac{e^{-\sqrt{p_y^2 + k_n^2}|x-x'|}}{|x-x'|}\exp\left\{i\frac{\beta}{\gamma}x'\cdot(p_{y_1} - p'_{y_1})\right\}\widehat{|V_{23}|^{1/2}}((p_{y_1} - p'_{y_1})/\gamma), \tag{38}$$

$\beta := -m_2\hbar/((m_1+m_2)\sqrt{2\mu_{12}})$ and $\gamma := \hbar/\sqrt{2M_{12}}$. Using the estimate

$$\|\mathcal{K}_n\|^2 \leq \sup_{p_{y_r}} \int d^3x\, d^3x'\, d^3p_{y_1}\, d^3p'_{y_1}\, |K_n(x, x', p_{y_1}, p'_{y_1}; p_{y_r})|^2 \tag{39}$$

we get

$$\|\mathcal{K}_n\|^2 \leq C_0 \sup_{|p_{y_r}| \leq \sqrt{1-p_{y_1}^2}} \int_{|p_{y_1}|\leq 1} d^3p_{y_1} \left[\frac{1}{k_n+\sqrt{|p_y|}} - \frac{1}{k_n+1}\right]^2 \frac{1}{\sqrt{p_y^2+k_n^2}}, \tag{40}$$

where the constant

$$C_0 := \frac{1}{2^7\pi^5\gamma^6}\left(\int d^3x\, (V_{12})_-(\alpha x)\right)\left(\int d^3s\, \left|\widehat{|V_{23}|^{1/2}}(s/\gamma)\right|^2\right)\left(\int d^3t\, |t|^{-2}e^{-2|t|}\right) \tag{41}$$

is finite. Continuing (40)

$$\|\mathcal{K}_n\|^2 \leq C_0 \int_{|p_{y_1}|\leq 1} d^3p_{y_1} \frac{1}{|p_{y_1}|^2} = 4\pi C_0. \tag{42}$$

In the case (b) we make the orthogonal transformation of Jacobi coordinates, where $\tilde{y}_1 = \alpha'^{-1}(r_4 - r_3)$ and $\alpha' := \hbar/\sqrt{2\mu_{34}}$. Other transformed coordinates we denote as $\tilde{y}_r := (\tilde{y}_2, \ldots, \tilde{y}_{N-2}) \in \mathbb{R}^{3N-9}$. This choice of coordinates for $N = 4$ is illustrated in Fig. 1 (Right). We need to prove that $\|\mathcal{L}_n\|$ is uniformly bounded, where

$$\mathcal{L}_n = \tilde{\mathcal{F}}_{12}\sqrt{(v_{12})_-}\Big[H_0 + k_n^2\Big]^{-1}\Big\{B_{12}^{-1}(k_n) - 1\Big\}\sqrt{|v_{34}|}\tilde{\mathcal{F}}_{12}^{-1} \tag{43}$$

and $\tilde{\mathcal{F}}_{12}$ is defined as in (9). The operator $\mathcal{L}_n$ acts on $\phi(x, \tilde{p}_{y_1}, \tilde{p}_{y_r}) \in L^2(\mathbb{R}^{3N-3})$ as

$$\mathcal{L}_n\phi(x, \tilde{p}_{y_1}, \tilde{p}_{y_r}) = \int d^3x'\, d^3p'_{y_1}\, L_n(x, x', \tilde{p}_{y_1}, \tilde{p}'_{y_1}; \tilde{p}_{y_r})\phi(x', \tilde{p}'_{y_1}, \tilde{p}_{y_r}), \tag{44}$$

where the integral kernel is

$$\begin{aligned} L_n(x, x', \tilde{p}_{y_1}, \tilde{p}'_{y_1}; \tilde{p}_{y_r}) &= \frac{1}{2^{7/2}\pi^{5/2}(\alpha')^3}\left\{\frac{1}{k_n+1+t(\tilde{p}_y)} - \frac{1}{k_n+1}\right\}\Big|(V_{12})_-(\alpha x)\Big|^{1/2} \\ &\quad\times \frac{e^{-\sqrt{\tilde{p}_y^2+k_n^2}|x-x'|}}{|x-x'|}\widehat{|V_{34}|^{1/2}}((\tilde{p}_{y_1} - \tilde{p}'_{y_1})/\alpha'). \end{aligned} \tag{45}$$

Now the proof that $\|\mathcal{L}_n\|$ is uniformly bounded is identical to the one in the case (a) and so we omit it. The general case of $D^{(2)}_{12;\delta}$ follows from (a) and (b) by making an orthogonal coordinate transformation, which corresponds to the appropriate permutation of the particle numbers. □

Acknowledgments

The author would like to thank Prof. Walter Greiner for the warm hospitality at FIAS.

Appendix A. The No-Clustering Theorem

Below we prove the statement, which we call the no–clustering theorem. In the following $\chi_L : \mathbb{R}^3 \to \mathbb{R}$ denotes the function such that $\chi_L(r) = 1$ if $0 \leq |r| \leq L$ and zero otherwise.

We shall make use of the following Lemma concerning minimizing sequences [16].

Lemma 3 ((Zhislin)). *Suppose that* $H \geq 0$ *is given by (1), where* $\lambda = 1$ *and* $V_{ij} \in L^2 + L^\infty_\infty$. *Suppose additionally that there is a normalized minimizing sequence* $f_n \in D(H_0)$ *such that* $(f_n, Hf_n) \to 0$. *If* f_n *does not totally spread then there exists a normalized* $\phi_0 \in D(H_0)$ *such that* $H\phi_0 = 0$.

Proof. Since f_n does not totally spread there must exist a subsequence such that $\|\chi_{\{x||x|\leq R\}} f_{n_k}\| > a$ for some $R > 0$ and $a > 0$. We can assume that $f_{n_k} \overset{w}{\to} \phi \in L^2(\mathbb{R}^{3N-3})$ otherwise we could pass to the weakly converging subsequence, which exists by the Banach–Alaoglu theorem. Thus for any $g \in D(H_0)$ we have

$$(Hg, \phi) = \lim_{k\to\infty} (Hg, f_{n_k}) = \lim_{k\to\infty} (H^{1/2}g, H^{1/2}f_{n_k}) = 0 \tag{A.1}$$

because $\|H^{1/2}f_{n_k}\| = (f_{n_k}, Hf_{n_k}) \to 0$ by condition of the lemma. From (A.1) it follows that $\phi \in D(H_0)$ and $H\phi = 0$. That $\|\phi\| \neq 0$ follows from Lemma 3 in [1]. Setting $\phi_0 = \phi/\|\phi\|$ we prove the lemma. □

Theorem 3. *Suppose that* H *is given by (1), where* $\lambda = 1$ *and* $V_{ij} \in L^2 + L^\infty_\infty$, *and none of the subsystems has an eigenstate with an energy less or equal to zero. Let* $\psi_n \in D(H_0)$ *be a totally spreading sequence such that* $(\psi_n, H\psi_n) \to 0$. *Then* $(\psi_n, F(r_i - r_j)\psi_n) \to 0$ *for all particle pairs* (i, j) *and any given* $F \in L^2(\mathbb{R}^3) + L^\infty_\infty(\mathbb{R}^3)$.

Proof. Note that $\|H_0\psi_n\|$ is uniformly bounded, c. f. Lemma 1 in [1] and $\psi_n \overset{w}{\to} 0$ because ψ_n totally spreads. By Lemma 4 it is enough to prove the statement for $F(r) = \chi_L(r)$ and all $L > 0$. For $N = 2$ the statement becomes trivial. For $N \geq 3$ we prove the theorem by induction assuming that it holds for $N - 1$ particles. Without loosing generality it is enough to show that $(\psi_n, \chi_L(r_1 - r_2)\psi_n) \to 0$ for all $L > 0$.

We can assume that $\psi_n \in C_0^\infty(\mathbb{R}^{3N-3})$ otherwise we can pass to an appropriate sequence using that $C_0^\infty(\mathbb{R}^{3N-3})$ is dense in $D(H_0)$, see [17]. A proof by contradiction. Lets us assume that

$$\limsup_{n\to\infty} (\psi_n, \chi_L(r_1 - r_2)\psi_n) = a' \tag{A.2}$$

for some $L > 0$ and $a' > 0$.

Let $J_s \in C^2(\mathbb{R}^{3N-3})$ denote the Ruelle–Simon partition of unity, see Definition 3.4 and Proposition 3.5 in [10]. For $s = 1, 2, \ldots, N$ one has $J_s \geq 0$, $\sum_s J_s^2 = 1$ and $J_s(\lambda x) = J_s(x)$ for $\lambda \geq 1$ and $|x| = 1$. Besides there exists $C > 0$ such that for $i \neq s$

$$\operatorname{supp} J_s \cap \{x||x| > 1\} \subset \{x|\ |r_i - r_s| \geq C|x|\}. \tag{A.3}$$

By the IMS formula (Theorem 3.2 in [10])

$$H = \sum_{s=1}^{N} J_s H_s J_s + K, \tag{A.4}$$

where

$$K := \sum_{s} \sum_{l \neq s} V_{ls} |J_s|^2 + \sum_{s} |\nabla J_s|^2, \tag{A.5}$$

$$H_s := H - \sum_{l \neq s} V_{ls}. \tag{A.6}$$

The operator K is relatively H_0–compact, see Lemma 7.11 in [18]. H_s is the same operator as H except that the pair–interactions that involve particle s are switched off. By (A.2) we get

$$\limsup_{n\to\infty} \sum_{s} (\psi_n^{(s)}, \chi_L(r_1 - r_2)\psi_n^{(s)}) = a', \tag{A.7}$$

where we define $\psi_n^{(s)} := J_s \psi_n \in C_0^\infty(\mathbb{R}^{3N-3})$. The operators $J_1\chi_L(r_1 - r_2)$ and $J_2\chi_L(r_1 - r_2)$ are relatively H_0 compact, hence, $\|\chi_L(r_1 - r_2)\psi_n^{(1)}\| \to 0$ and $\|\chi_L(r_1 - r_2)\psi_n^{(2)}\| \to 0$ by Lemma 2 in [1]. Thus there must exist $s_0 \geq 3$ such that

$$\limsup_{n\to\infty} (\psi_n^{(s_0)}, \chi_L(r_1 - r_2)\psi_n^{(s_0)}) = 2a, \tag{A.8}$$

where $0 < a \leq 1/2$ is a constant. Let $\zeta \in \mathbb{R}^{3N-6}$ denote the internal Jacobi coordinates for the particles $\{1, 2, \ldots s_0 - 1, s_0 + 1, \ldots, N\}$ and $y \in \mathbb{R}^3$ the coordinate, which points from the particle s_0 to the center of mass of other particles. We choose the scales so that $H_0 = -\Delta_\zeta - \Delta_y$. It is convenient to introduce

$$\begin{aligned} H^{(s_0)} &:= -\Delta_\zeta + V(\zeta), \\ V(\zeta) &:= \sum_{i<k} V_{ik} - \sum_{l \neq s_0} V_{ls_0}. \end{aligned} \tag{A.9}$$

Clearly,

$$H_{s_0} \geq H^{(s_0)} \geq 0. \tag{A.10}$$

The operator $H^{(s_0)}$ is the Hamiltonian of the particles $\{1, 2, \ldots s_0 - 1, s_0 + 1, \ldots, N\}$ and can be considered on the domain $D(-\Delta_\zeta) \subset L^2(\mathbb{R}^{3N-6})$ as well. We have $\psi_n \xrightarrow{w} 0$ because ψ_n totally spreads. Because K in (A.5) is relatively H_0 compact we have $K\psi_n \to 0$, see Lemma 2 in [1]. Using $(\psi_n, H\psi_n) \to 0$ and $H_s \geq 0$ we infer from (A.4) that $(\psi_n^{(s)}, H_s\psi_n^{(s)}) \to 0$ for all s. Hence, by (A.10)

$$(\psi_n^{(s_0)}, H^{(s_0)}\psi_n^{(s_0)}) \to 0. \tag{A.11}$$

Looking at (A.8) and (A.11) we conclude that there exists a subsequence $\psi_{n_k}^{(s_0)}$ such that

$$(\psi_{n_k}^{(s_0)}, \chi_L(r_1 - r_2)\psi_{n_k}^{(s_0)}) \geq a, \tag{A.12}$$

$$(\psi_{n_k}^{(s_0)}, H^{(s_0)}\psi_{n_k}^{(s_0)}) \to 0. \tag{A.13}$$

From (A.12) it follows that $\sqrt{a} \leq \|\psi^{(s_0)}_{n_k}\| \leq 1$. Thus defining $g_k := \psi^{(s_0)}_{n_k}/\|\psi^{(s_0)}_{n_k}\|$ we obtain

$$(g_k, \chi_L(r_1 - r_2)g_k) \geq a \tag{A.14}$$

$$\varepsilon_k := (g_k, H^{(s_0)}g_k) \to 0, \tag{A.15}$$

where $g_k \in C_0^\infty(\mathbb{R}^{3N-3})$ and $\|g_k\| = 1$. For $f(\zeta, y), h(\zeta, y) \in L^2(\mathbb{R}^{3N-3})$ let us introduce the notation

$$(f, h)_\zeta := \int d^{3N-6}\zeta \; f^*(\zeta, y)h(\zeta, y), \tag{A.16}$$

where $(f, h)_\zeta$ depends on $y \in \mathbb{R}^3$.

Now we define the following subsets of $\mathbb{R}^3$

$$\mathcal{M}_k := \left\{ y \middle| (g_k, g_k)_\zeta > 0 \right\} \cap \left\{ y \middle| (g_k, H^{(s_0)}g_k)_\zeta < \sqrt{\varepsilon_k}(g_k, g_k)_\zeta \right\}, \tag{A.17}$$

$$\mathcal{N}_k := \left\{ y \middle| (g_k, \chi_L(r_1 - r_2)g_k)_\zeta \geq (a/2)(g_k, g_k)_\zeta \right\}. \tag{A.18}$$

By standard results $\mathcal{M}_k, \mathcal{N}_k$ are Borel sets. Below we prove that there exists k_0 such that $\mathcal{N}_k \cap \mathcal{M}_k \neq \emptyset$ for $k \geq k_0$. For any Borel set $X \subset \mathbb{R}^3$ we define

$$\mu_k(X) := \int_X d^3y \; (g_k, g_k)_\zeta. \tag{A.19}$$

Because g_k is normalized we have $\mu_k(\mathbb{R}^3) = 1$. On one hand, using (A.15) and (A.17)

$$\mu_k(\mathbb{R}^3/\mathcal{M}_k) = \int_{\mathbb{R}^3/\mathcal{M}_k} d^3y \; (g_k, g_k)_\zeta \leq \frac{1}{\sqrt{\varepsilon_k}} \int_{\mathbb{R}^3/\mathcal{M}_k} d^3y \; (g_k, H^{(s_0)}g_k)_\zeta \leq \sqrt{\varepsilon_k}. \tag{A.20}$$

Hence,

$$\mu_k(\mathcal{M}_k) \geq 1 - \sqrt{\varepsilon_k}. \tag{A.21}$$

On the other hand, using that according to (A.14) $\int d^3y \; (g_k, \chi_L(r_1 - r_2)g_k)_\zeta \geq a$ we get

$$\begin{aligned} \mu_k(\mathcal{N}_k) &\geq \int_{\mathcal{N}_k} d^3y (g_k, \chi_L(r_1 - r_2)g_k)_\zeta \geq a - \int_{\mathbb{R}^3/\mathcal{N}_k} (g_k, \chi_L(r_1 - r_2)g_k)_\zeta \\ &\geq a - \frac{a}{2}\mu_k(\mathbb{R}^3/\mathcal{N}_k) \geq \frac{a}{2}, \end{aligned} \tag{A.22}$$

where we applied (A.18) and $\mu_k(\mathbb{R}^m/\mathcal{N}_k) \leq 1$. Now it is clear that that there exists k_0 such that $\mathcal{N}_k \cap \mathcal{M}_k \neq \emptyset$ for $k \geq k_0$. Otherwise, according to (A.21) and (A.22) we would have

$$1 = \mu_k(\mathbb{R}^m) \geq \mu_k(\mathcal{M}_k) + \mu_k(\mathcal{N}_k) \geq 1 + \frac{a}{2} - \sqrt{\varepsilon_k}, \tag{A.23}$$

which is a contradiction since $\varepsilon_k \to 0$. Now we construct the minimizing sequence for $H^{(s_0)}$ (considered now on $D(-\Delta_\zeta)$) taking any $y_k \in \mathcal{N}_k \cap \mathcal{M}_k$ for $k \geq k_0$ and setting

$$\phi_k(\zeta) := g_k(y_k, \zeta) \left(\int d^{3N-6}\zeta \; |g_k(y_k, \zeta)| \right)^{-1/2}. \tag{A.24}$$

Due to (A.17)–(A.18) the sequence $\phi_k(\zeta) \in C_0^\infty(\mathbb{R}^{3N-6})$ has the following properties: $\|\phi_k\| = 1$, $(\phi_k, H^{(s_0)}\phi_k) \to 0$ and

$$(\phi_k, \chi_L(r_1 - r_2)\phi_k) \geq a/2. \tag{A.25}$$

By Lemma 3 ϕ_k must totally spread because $H^{(s_0)}$ is not allowed to have zero energy bound states. Since $H^{(s_0)}$ is the Hamiltonian of $N-1$ particles by the induction assumption it follows that

$$(\phi_k, \chi_L(r_1 - r_2)\phi_k) \to 0, \tag{A.26}$$

which contradicts (A.25). □

Lemma 4. *Let $f_n(x) \in D(H_0) \subset L^2(\mathbb{R}^{3N-3})$ and $\|f_n\| + \|H_0 f_n\| \leq 1$. Suppose that $\|\chi_{\{x||r_i-r_j|<q\}} f_n\| \to 0$ for some fixed $i \neq j$ and any $q > 0$. Then $(f_n, F(r_i - r_j)f_n) \to 0$ for any given $F \in L^2(\mathbb{R}^3) + L^\infty_\infty(\mathbb{R}^3)$.*

Proof. Obviously, it suffices to consider $F \in L^2(\mathbb{R}^3)$.

$$\||F|^{1/2} f_n\| \leq (\chi_{\{x||r_i-r_j|<q\}} f_n, |F| f_n) + (\chi_{\{x||r_i-r_j|\geq q\}} f_n, |F| f_n) \tag{A.27}$$

$$= (\chi_{\{x||r_i-r_j|<q\}} f_n, |F| f_n) + (f_n, \chi_{\{x||r_i-r_j|\geq q\}} |F|(H_0+1)^{-1}(H_0+1) f_n) \tag{A.28}$$

$$\leq \|\chi_{\{x||r_i-r_j|<q\}} f_n\| \, \||F| f_n\| + \Big\|\chi_{\{x||r_i-r_j|\geq q\}} |F|(H_0+1)^{-1}\Big\|. \tag{A.29}$$

The first term in (A.29) goes to zero because $|F(r_i - r_j)|$ is relatively H_0 bounded. The second term is an operator norm, which can be made as small as pleased by setting q large enough, see Lemma 5 in [1]. □

Acknowledgments

The author would like to thank Prof. Walter Greiner for the warm hospitality at FIAS.

References

[1] D. K. Gridnev, *Zero Energy Bound States and Resonances in Three–Particle Systems*, arXiv:1111.6788v1

[2] J. M. Richard, Phys. Rev. **A 67**, 034702 (2003)

[3] J. M. Richard and S. Fleck, Phys. Rev. Lett. 73, 1464 (1994)

[4] M. Klaus and B. Simon, Ann. Phys. (N.Y.) **130**, 251 (1980)

[5] M. Klaus and B. Simon, Comm. Math. Phys. **78**, 153 (1980)

[6] F. Ahia, J. Math. Phys. **33**, 189 (1992)

[7] A. V. Sobolev, Commun. Math. Phys. **156**, 101 (1993)

[8] D. R. Yafaev, Math. USSR-Sb. **23**, 535-559 (1974); Notes of LOMI Seminars **51** (1975) (Russian)

[9] R. Courant and D. Hilbert, *Methods of Mathematical Physics*, Interscience, New York, 1953, Vol. 1 p. 446

[10] H. L. Cycon, R. G. Froese, W. Kirsch and B. Simon, *Schrödinger Operators with Applications to Quantum Mechanics and Global Geometry*, Springer–Verlag, Berlin Heidelberg (1987)

[11] M. Reed and B. Simon, *Methods of Modern Mathematical Physics*, vol. 2 Academic Press/New York (1975) and vol. 4, Academic Press/New York (1978).

[12] W. Greiner, *Quantum Mechanics: An Introduction*, Springer–Verlag, Berlin (2000)

[13] B. Simon, Bull. Amer. Math. Soc. **7** 447 (1982)

[14] F. Gesztesy and B. Simon, Comm. Math. Phys. **161** 503 (1988); Trans. Amer. Math. Soc. **335** 329 (1993)

[15] B. Simon, *Trace Ideals and Their Applications*, Cambridge University Press, 1979

[16] G. M. Zhislin, Trudy Mosk. Mat. Obšč. **9**, 81 (1960); E. F. Zhizhenkova and G. M. Zhislin, Trudy Mosk. Mat. Obšč. **9**, 121 (1960)

[17] E. H. Lieb and M. Loss, *Analysis*, American Mathematical Society (1997)

[18] G. Teschl, *Mathematical Methods in Quantum Mechanics; With Applications to Schrödinger Operators*, Lecture Notes (2005), http://www.mat.univie.ac.at/∼gerald/ftp/book-schroe/index.html